

NOTICE

The following two-volume report is intended solely as guidance to EPA and other environmental professionals. This document does not constitute rulemaking by the Agency, and cannot be relied on to create a substantive or procedural right enforceable by any party in litigation with the United States. EPA may take action that is at variance with the information, policies, and procedures in this document and may change them at any time without public notice.

Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government.

FOREWORD

Understanding the long-term behavior of contaminants in the subsurface is becoming increasingly more important as the nation addresses groundwater contamination. Groundwater contamination is a national concern as about 50 percent of the United States population receives its drinking water from groundwater. It is the goal of the Environmental Protection Agency (EPA) to prevent adverse effects to human health and the environment and to protect the environmental integrity of the nation's groundwater.

Once groundwater is contaminated, it is important to understand how the contaminant moves in the subsurface environment. Proper understanding of the contaminant fate and transport is necessary in order to characterize the risks associated with the contamination and to develop, when necessary, emergency or remedial action plans. The parameter known as the partition (or distribution) coefficient (K_d) is one of the most important parameters used in estimating the migration potential of contaminants present in aqueous solutions in contact with surface, subsurface and suspended solids.

This two-volume report describes: (1) the conceptualization, measurement, and use of the partition coefficient parameter; and (2) the geochemical aqueous solution and sorbent properties that are most important in controlling adsorption/retardation behavior of selected contaminants. Volume I of this document focuses on providing EPA and other environmental remediation professionals with a reasoned and documented discussion of the major issues related to the selection and measurement of the partition coefficient for a select group of contaminants. The selected contaminants investigated in this two-volume document include: chromium, cadmium, cesium, lead, plutonium, radon, strontium, thorium, tritium (^3H), and uranium. This two-volume report also addresses a void that has existed on this subject in both this Agency and in the user community.

It is important to note that soil scientists and geochemists knowledgeable of sorption processes in natural environments have long known that generic or default partition coefficient values found in the literature can result in significant errors when used to predict the absolute impacts of contaminant migration or site-remediation options. Accordingly, one of the major recommendations of this report is that for site-specific calculations, partition coefficient values measured at site-specific conditions are absolutely essential.

For those cases when the partition coefficient parameter is not or cannot be measured, Volume II of this document: (1) provides a "thumb-nail sketch" of the key geochemical processes affecting the sorption of the selected contaminants; (2) provides references to related key experimental and review articles for further reading; (3) identifies the important aqueous- and solid-phase parameters controlling the sorption of these contaminants in the subsurface environment under oxidizing conditions; and (4) identifies, when possible, minimum and maximum conservative partition coefficient values for each contaminant as a function of the key geochemical processes affecting their sorption.

This publication is the result of a cooperative effort between the EPA Office of Radiation and Indoor Air, Office of Solid Waste and Emergency Response, and the Department of Energy Office of Environmental Restoration (EM-40). In addition, this publication is produced as part of ORIA's long-term strategic plan to assist in the remediation of contaminated sites. It is published and made available to assist all environmental remediation professionals in the cleanup of groundwater sources all over the United States.

Stephen D. Page, Director
Office of Radiation and Indoor Air

ACKNOWLEDGMENTS

Ronald G. Wilhelm from ORIA's Center for Remediation Technology and Tools was the project lead and EPA Project Officer for this two-volume report. Paul Beam, Environmental Restoration Program (EM-40), was the project lead and sponsor for the Department of Energy (DOE). Project support was provided by both DOE/EM-40 and EPA's Office of Remedial and Emergency Response (OERR).

EPA/ORIA wishes to thank the following people for their assistance and technical review comments on various drafts of this report:

Patrick V. Brady, U.S. DOE, Sandia National Laboratories
David S. Brown, U.S. EPA, National Exposure Research Laboratory
Joe Eidelberg, U.S. EPA, Region 9
Amy Gamerdinger, Washington State University
Richard Graham, U.S. EPA, Region 8
John Griggs, U.S. EPA, National Air and Radiation Environmental Laboratory
David M. Kargbo, U.S. EPA, Region 3
Ralph Ludwig, U.S. EPA, National Risk Management Research Laboratory
Irma McKnight, U.S. EPA, Office of Radiation and Indoor Air
William N. O'Steen, U.S. EPA, Region 4
David J. Reisman, U.S. EPA, National Risk Management Research Laboratory
Kyle Rogers, U.S. EPA, Region 5
Joe R. Williams, U.S. EPA, National Risk Management Research Laboratory
OSWER Regional Groundwater Forum Members

In addition, special acknowledgment goes to Carey A. Johnston from ORIA's Center for Remediation Technology and Tools for his contributions in the development, production, and review of this document.

Principal authorship in production of this guide was provided by the Department of Energy's Pacific Northwest National Laboratory (PNNL) under the Interagency Agreement Number DW89937220-01-03. Lynnette Downing served as the Department of Energy's Project Officer for this Interagency Agreement. PNNL authors involved in this project include:

Kenneth M. Krupka
Daniel I. Kaplan
Gene Whelan
R. Jeffrey Serne
Shas V. Mattigod

**TO COMMENT ON THIS GUIDE OR PROVIDE INFORMATION FOR FUTURE
UPDATES:**

Send all comments/updates to:

U.S. Environmental Protection Agency
Office of Radiation and Indoor Air
Attention: Understanding Variation in Partition (K_d) Values
401 M Street, SW (6602J)
Washington, DC 20460

or

radiation.questions@epa.gov

ABSTRACT

This two-volume report describes the conceptualization, measurement, and use of the partition (or distribution) coefficient, K_d , parameter, and the geochemical aqueous solution and sorbent properties that are most important in controlling adsorption/retardation behavior of selected contaminants. The report is provided for technical staff from EPA and other organizations who are responsible for prioritizing site remediation and waste management decisions. Volume I discusses the technical issues associated with the measurement of K_d values and its use in formulating the retardation factor, R_f . The K_d concept and methods for measurement of K_d values are discussed in detail in Volume I. Particular attention is directed at providing an understanding of: (1) the use of K_d values in formulating R_f , (2) the difference between the original thermodynamic K_d parameter derived from ion-exchange literature and its “empiricized” use in contaminant transport codes, and (3) the explicit and implicit assumptions underlying the use of the K_d parameter in contaminant transport codes. A conceptual overview of chemical reaction models and their use in addressing technical defensibility issues associated with data from K_d studies is presented. The capabilities of EPA’s geochemical reaction model MINTEQA2 and its different conceptual adsorption models are also reviewed. Volume II provides a “thumb-nail sketch” of the key geochemical processes affecting the sorption of selected inorganic contaminants, and a summary of K_d values given in the literature for these contaminants under oxidizing conditions. The contaminants chosen for the first phase of this project include chromium, cadmium, cesium, lead, plutonium, radon, strontium, thorium, tritium (^3H), and uranium. Important aqueous speciation, (co)precipitation/dissolution, and adsorption reactions are discussed for each contaminant. References to related key experimental and review articles for further reading are also listed.

CONTENTS

	<u>Page</u>
NOTICE	ii
FOREWORD	iii
ACKNOWLEDGMENTS	v
FUTURE UPDATES	vi
ABSTRACT	vii
LIST OF FIGURES	xiii
LIST OF TABLES	xv
1.0 Introduction	1.1
2.0 The K_d Model	2.1
3.0 Methods, Issues, and Criteria for Measuring K_d Values	3.1
3.1 Laboratory Batch Methods	3.1
3.2 Laboratory Flow-Through Method	3.1
3.3 Other Methods	3.2
3.4 Issues	3.2
4.0 Application of Chemical Reaction Models	4.1
5.0 Contaminant Geochemistry and K_d Values	5.1
5.1 General	5.1
5.2 Cadmium Geochemistry and K_d Values	5.5
5.2.1 Overview: Important Aqueous- and Solid-Phase Parameters Controlling Retardation	5.5
5.2.2 General Geochemistry	5.5
5.2.3 Aqueous Speciation	5.6
5.2.4 Dissolution/Precipitation/Coprecipitation	5.8
5.2.5 Sorption/Desorption	5.9
5.2.6 Partition Coefficient, K_d , Values	5.10
5.2.6.1 General Availability of K_d Values	5.10
5.2.6.2 Look-Up Tables	5.11
5.2.6.2.1 Limits of K_d Values with Aluminum/Iron-Oxide Concentrations	5.11
5.2.6.2.2 Limits of K_d Values with Respect to CEC	5.12
5.2.6.2.3 Limits of K_d Values with Respect to Clay Concentrations	5.12
5.2.6.2.4 Limits of K_d Values with Respect to Concentration of Organic Matter	5.12
5.2.6.2.5 Limits of K_d Values with Respect to Dissolved Calcium,	

Magnesium, and Sulfide Concentrations, and Redox Conditions	5.12
5.3 Cesium Geochemistry and K_d Values	5.13
5.3.1 Overview: Important Aqueous- and Solid-Phase Parameters Controlling Retardation	5.13
5.3.2 General Geochemistry	5.13
5.3.3 Aqueous Speciation	5.13
5.3.4 Dissolution/Precipitation/Coprecipitation	5.14
5.3.5 Sorption/Desorption	5.14
5.3.6 Partition Coefficient, K_d , Values	5.15
5.3.6.1 General Availability of K_d Data	5.15
5.3.6.2 Look-Up Tables	5.16
5.3.6.2.1 Limits of K_d with Respect to pH	5.18
5.3.6.2.2 Limits of K_d with Respect to Potassium, Ammonium, and Aluminum/Iron-Oxide Concentrations	5.18
5.4 Chromium Geochemistry and K_d Values	5.18
5.4.1 Overview: Important Aqueous- and Solid-Phase Parameters Controlling Retardation	5.18
5.4.2 General Geochemistry	5.18
5.4.3 Aqueous Speciation	5.19
5.4.4 Dissolution/Precipitation/Coprecipitation	5.19
5.4.5 Sorption/Desorption	5.20
5.4.6 Partition Coefficient, K_d , Values	5.21
5.4.6.1 General Availability of K_d Data	5.21
5.4.6.2 Look-Up Tables	5.22
5.4.6.2.1 Limits of K_d with Respect to pH	5.23
5.4.6.2.2 Limits of K_d with Respect to Extractable Iron Content	5.23
5.4.6.2.3 Limits of K_d with Respect to Competing Anion Concentrations . .	5.23
5.5 Lead Geochemistry and K_d Values	5.25
5.5.1 Overview: Important Aqueous- and Solid-Phase Parameters Controlling Retardation	5.25
5.5.2 General Geochemistry	5.25
5.5.3 Aqueous Speciation	5.26
5.5.4 Dissolution/Precipitation/Coprecipitation	5.27
5.5.5 Sorption/Desorption	5.30
5.5.6 Partition Coefficient, K_d , Values	5.31
5.5.6.1 General Availability of K_d Data	5.31
5.5.6.2 K_d Look-Up Tables	5.33
5.5.6.2.1 Limits of K_d with Respect to pH	5.33
5.5.6.2.2 Limits of K_d with Respect to Equilibrium Lead Concentrations Extractable Iron Content	5.34

5.6 Plutonium Geochemistry and K_d Values	5.34
5.6.1 Overview: Important Aqueous- and Solid-Phase Parameters	
Controlling Retardation	5.34
5.6.2 General Geochemistry	5.34
5.6.3 Aqueous Speciation	5.35
5.6.4 Dissolution/Precipitation/Coprecipitation	5.37
5.6.5 Sorption/Desorption	5.40
5.6.6 Partition Coefficient, K_d , Values	5.41
5.6.6.1 General Availability of K_d Data	5.41
5.6.6.2 K_d Look-Up Tables	5.43
5.6.6.2.1 Limits of K_d with Respect to Clay Content	5.43
5.6.6.2.2 Limits of K_d with Respect to Dissolved Carbonate	
Concentrations	5.44
5.7 Radon Geochemistry and K_d Values	5.44
5.7.1 Overview: Important Aqueous- and Solid-Phase Parameters	
Controlling Retardation	5.44
5.7.2 General Geochemistry	5.45
5.7.3 Aqueous Speciation	5.45
5.7.4 Dissolution/Precipitation/Coprecipitation	5.46
5.7.5 Sorption/Desorption	5.46
5.7.6 Partition Coefficient, K_d , Values	5.46
5.8 Strontium Geochemistry and K_d Values	5.46
5.8.1 Overview: Important Aqueous- and Solid-Phase Parameters	
Controlling Retardation	5.46
5.8.2 General Geochemistry	5.47
5.8.3 Aqueous Speciation	5.47
5.8.4 Dissolution/Precipitation/Coprecipitation	5.48
5.8.5 Sorption/Desorption	5.49
5.8.6 Partition Coefficient, K_d , Values	5.51
5.8.6.1 General Availability of K_d Data	5.51
5.8.6.2 Look-Up Tables	5.51
5.8.6.2.1 Limits of K_d with Respect to pH, CEC, and	
Clay Concentrations Values	5.52
5.8.6.2.2 Limits of K_d with Respect to Dissolved Calcium	
Concentrations	5.52
5.8.6.2.3 Limits of K_d with Respect to Dissolved Stable	
Strontium and Carbonate Concentrations	5.53
5.9 Thorium Geochemistry and K_d Values	5.53
5.9.1 Overview: Important Aqueous- and Solid-Phase Parameters	
Controlling Retardation	5.53
5.9.2 General Geochemistry	5.54

5.9.3 Aqueous Speciation	5.55
5.9.4 Dissolution/Precipitation/Coprecipitation	5.58
5.9.5 Sorption/Desorption	5.60
5.9.6 Partition Coefficient, K_d , Values	5.61
5.9.6.1 General Availability of K_d Data	5.61
5.9.6.2 Look-Up Tables	5.62
5.9.6.2.1 Limits of K_d with Respect to Organic Matter and Aluminum/Iron-Oxide Concentrations	5.63
5.9.6.2.2 Limits of K_d with Respect to Dissolved Carbonate Concentrations	5.63
5.10 Tritium Geochemistry and K_d Values	5.64
5.10.1 Overview: Important Aqueous- and Solid-Phase Parameters Controlling Retardation	5.64
5.10.2 General Geochemistry	5.64
5.10.3 Aqueous Speciation	5.65
5.10.4 Dissolution/Precipitation/Coprecipitation	5.65
5.10.5 Sorption/Desorption	5.65
5.10.6 Partition Coefficient, K_d , Values	5.65
5.11 Uranium Geochemistry and K_d Values	5.65
5.11.1 Overview: Important Aqueous- and Solid-Phase Parameters Controlling Retardation	5.65
5.11.2 General Geochemistry	5.66
5.11.3 Aqueous Speciation	5.67
5.11.4 Dissolution/Precipitation/Coprecipitation	5.69
5.11.5 Sorption/Desorption	5.72
5.11.6 Partition Coefficient, K_d , Values	5.74
5.11.6.1 General Availability of K_d Data	5.74
5.11.6.2 Look-Up Table	5.74
5.11.6.2.1 Limits K_d Values with Respect to Dissolved Carbonate Concentrations	5.75
5.11.6.2.2 Limits of K_d Values with Respect to Clay Content and CEC ...	5.76
5.11.6.2.3 Use of Surface Complexation Models to Predict Uranium K_d Values	5.76
5.12 Conclusions	5.77
6.0 References	6.1
Appendix A - Acronyms and Abbreviations	A.1
Appendix B - Definitions	B.1

Appendix C - Partition Coefficients for Cadmium	C.1
Appendix D - Partition Coefficients for Cesium	D.1
Appendix E - Partition Coefficients for Chromium	E.1
Appendix F - Partition Coefficients for Lead	F.1
Appendix G - Partition Coefficients for Plutonium	G.1
Appendix H - Partition Coefficients for Strontium	H.1
Appendix I - Partition Coefficients for Thorium	I.1
Appendix J - Partition Coefficients for Uranium	J.1

LIST OF FIGURES

	<u>Page</u>
Figure 5.1. Calculated distribution of cadmium aqueous species as a function of pH for the water composition in Table 5.1	5.7
Figure 5.2. Calculated distribution of lead aqueous species as a function of pH for the water composition listed in Table 5.1	5.29
Figure 5.3. Calculated distribution of plutonium aqueous species as a function of pH for the water composition in Table 5.1.	5.39
Figure 5.4. Calculated distribution of thorium hydrolytic species as a function of pH. ...	5.57
Figure 5.5. Calculated distribution of thorium aqueous species as a function of pH for the water composition in Table 5.1.	5.59
Figure 5.6a. Calculated distribution of U(VI) hydrolytic species as a function of pH at 0.1 µg/l total dissolved U(VI)	5.70
Figure 5.6b. Calculated distribution of U(VI) hydrolytic species as a function of pH at 1,000 µg/l total dissolved U(VI)	5.71
Figure 5.7. Calculated distribution of U(VI) aqueous species as a function of pH for the water composition in Table 5.1	5.72
Figure C.1. Relation between cadmium K_d values and pH in soils	C.5
Figure D.1. Relation between cesium K_d values and CEC	D.7
Figure D.2. Relation between CEC and clay content	D.8
Figure D.3. K_d values calculated from an overall literature Freundlich equation for cesium (Equation D.2)	D.12
Figure D.4. Generalized cesium Freundlich equation (Equation D.3) derived from the literature	D.16
Figure D.5. Cesium K_d values calculated from generalized Freundlich equation (Equations D.3 and D.4) derived from the literature	D.16

Figure E.1.	Variation of K_d for Cr(VI) as a function of pH and DCB extractable Iron content without the presence of competing anions	E.10
Figure F.1.	Correlative relationship between K_d and pH	F.6
Figure F.2.	Variation of K_d as a function of pH and the equilibrium lead concentrations.	F.7
Figure G.1.	Scatter plot matrix of soil properties and the partition coefficient (K_d) of plutonium	G.12
Figure G.2.	Variation of K_d for plutonium as a function of clay content and dissolved carbonate concentrations.	G.14
Figure H.1.	Relation between strontium K_d values and CEC in soils.	H.5
Figure H.2.	Relation between strontium K_d values for soils with CEC values less than 15 meq/100 g.	H.7
Figure H.3.	Relation between strontium K_d values and soil clay content	H.7
Figure H.4.	Relation between strontium K_d values and soil pH	H.9
Figure I.1.	Linear regression between thorium K_d values and pH for the pH range from 4 to 8	I.5
Figure I.2.	Linear regression between thorium K_d values and pH for the pH range from 4 to 8	I.8
Figure J.1.	Field-derived K_d values for ^{238}U and ^{235}U from Serkiz and Johnson (1994) plotted as a function of porewater pH for contaminated soil/porewater samples	J.8
Figure J.2.	Field-derived K_d values for ^{238}U and ^{235}U from Serkiz and Johnson (1994) plotted as a function of the weight percent of clay-size particles in the contaminated soil/porewater samples	J.9
Figure J.3.	Field-derived K_d values for ^{238}U and ^{235}U plotted from Serkiz and Johnson (1994) as a function of CEC (meq/kg) of the contaminated soil/porewater samples	J.10
Figure J.4.	Uranium K_d values used for development of K_d look-up table	J.19

LIST OF TABLES

	<u>Page</u>
Table 5.1. Estimated mean composition of river water of the world from Hem (1985)	5.3
Table 5.2. Concentrations of contaminants used in the aqueous species distribution calculations.	5.4
Table 5.3. Cadmium aqueous species included in the speciation calculations	5.6
Table 5.4. Estimated range of K_d values for cadmium as a function of pH.	5.11
Table 5.5. Estimated range of K_d values (ml/g) for cesium based on CEC or clay content for systems containing <5 percent mica-like minerals in clay-size fraction and $<10^{-9}$ M aqueous cesium	5.17
Table 5.6. Estimated range of K_d values (ml/g) for cesium based on CEC or clay content for systems containing >5 percent mica-like minerals in clay-size fraction and $<10^{-9}$ M aqueous cesium	5.17
Table 5.7. Estimated range of K_d values for chromium (VI) as a function of soil pH, extractable iron content, and soluble sulfate	5.24
Table 5.8. Lead aqueous species included in the speciation calculations	5.28
Table 5.9. Estimated range of K_d values for lead as a function of soil pH, and equilibrium lead concentrations.	5.33
Table 5.10. Plutonium aqueous species included in the speciation calculations	5.38
Table 5.11. Estimated range of K_d values for plutonium as a function of the soluble carbonate and soil clay content values	5.43
Table 5.12. Strontium aqueous species included in the speciation calculations.	5.48
Table 5.13. Look-up table for estimated range of K_d values for strontium based on CEC (meq/100 g), clay content (wt.%), and pH.	5.53
Table 5.14. Thorium aqueous species included in the speciation calculations	5.56
Table 5.15. Look-up table for thorium K_d values (ml/g) based on pH and dissolved thorium concentrations.	5.63

Table 5.16. Uranium(VI) aqueous species included in the speciation calculations.	5.69
Table 5.17. Look-up table for estimated range of K_d values for uranium based on pH	5.75
Table 5.18. Selected chemical and transport properties of the contaminants.	5.78
Table 5.19. Distribution of dominant contaminant species at 3 pH values for an oxidizing water described in Tables 5.1 and 5.2.	5.79
Table 5.20. Some of the more important aqueous- and solid-phase parameters affecting contaminant sorption	5.81
Table C.1. Descriptive statistics of the cadmium K_d data set for soils	C.3
Table C.2. Correlation coefficients (r) of the cadmium K_d data set for soils	C.4
Table C.3. Look-up table for estimated range of K_d values for cadmium based on pH	C.5
Table C.4. Cadmium K_d data set for soils.	C.6
Table D.1. Descriptive statistics of cesium K_d data set including soil and pure mineral phases.	D.3
Table D.2. Descriptive statistics of data set including soils only.	D.4
Table D.3. Correlation coefficients (r) of the cesium K_d value data set that included soils and pure mineral phases.	D.6
Table D.4. Correlation coefficients (r) of the soil-only data set.	D.6
Table D.5. Effect of mineralogy on cesium exchange.	D.9
Table D.6. Cesium K_d values measured on mica (Fithian illite) via adsorption and desorption experiments.	D.10
Table D.7. Approximate upper limits of linear range of adsorption isotherms on various solid phases.	D.11
Table D.8. Freundlich equations identified in literature for cesium.	D.13
Table D.9. Descriptive statistics of the cesium Freundlich equations (Table D.8) reported in the literature.	D.15
Table D.10. Estimated range of K_d values (ml/g) for cesium based on CEC	

or clay content for systems <i>containing</i> <5% mica-like minerals in clay-size fraction and <10 ⁻⁹ M aqueous cesium	D.18
Table D.11. Estimated range of K _d values (ml/g) for cesium based on CEC or clay content for systems <i>containing</i> >5% mica-like minerals in clay-size fraction and <10 ⁻⁹ M aqueous cesium	D.18
Table D.12. Calculations for values used in look-up table.	D.19
Table D.13. Cesium K _d data base for soils and pure mineral phases	D.20
Table D.14. Cesium K _d data set for soils only	D.27
Table E.1. Summary of K _d values for Cr(VI) adsorption on soils.	E.5
Table E.2. Data from Rai et al. (1988) for the adsorption of Cr(VI) as a function of pH	E.8
Table E.3. Estimated range of K _d values for Cr(VI) as a function of soil pH, extractable iron content, and soluble sulfate.	E.9
Table E.4. Data from Rai et al. (1988) on effects of competing anions on Cr(VI) adsorption on Cecil/Pacolet soil.	E.11
Table E.5. Data from Rai et al. (1988) on effects of competing anions on Cr(VI) adsorption on Kenoma soil.	E.12
Table F.1. Summary of K _d values for lead adsorption on soils.	F.5
Table F.2. Estimated range of K _d values for lead as a function of soil pH, and equilibrium lead concentrations	F.8
Table G.1. Plutonium adsorption data for soil samples.	G.10
Table G.2. Regression models for plutonium adsorption.	G.13
Table G.3. Estimated range of K _d values for plutonium as a function of the soluble carbonate and soil clay content values.	G.13
Table H.1. Descriptive statistics of strontium K _d data set for soils.	H.3
Table H.2. Correlation coefficient (r) of the strontium K _d data set for soils	H.4
Table H.3. Simple and multiple regression analysis results involving strontium K _d values, CEC (meq/100 g), pH, and clay content (percent).	H.8

Table H.4. Look-up table for estimated range of K_d values for strontium based on CEC and pH.	H.10
Table H.5. Look-up table for estimated range of K_d values for strontium based on clay content and pH.	H.10
Table H.6. Calculations of clay content using regression equations containing CEC as a independent variable.	H.11
Table H.7. Strontium K_d data set for soils.	H.12
Table H.8. Strontium K_d data set for pure mineral phases and soils	H.16
Table I.1. Descriptive statistics of thorium K_d value data set presented in Section I.3.	I.3
Table I.2. Correlation coefficients (r) of the thorium K_d value data set presented in Section I.3.	I.4
Table I.3. Calculated aqueous speciation of thorium as a function of pH.	I.5
Table I.4. Regression coefficient and their statistics relating thorium K_d values and pH.	I.6
Table I.5. Look-up table for thorium K_d values (ml/g) based on pH and dissolved thorium concentrations.	I.7
Table I.6. Data set containing thorium K_d values.	I.9
Table J.1. Uranium K_d values (ml/g) listed by Warnecke et al. (1994, Table 1).	J.12
Table J.2. Uranium K_d values listed by McKinley and Scholtis (1993, Tables 1, 2, and 4) from sorption databases used by different international organizations for performance assessments of repositories for radioactive wastes.	J.17

Table J.3. Geometric mean uranium K_d values derived by Thibault et al. (1990) for sand, loam, clay, and organic soil types.	J.18
Table J.4. Look-up table for estimated range of K_d values for uranium based on pH.	J.22
Table J.5. Uranium K_d values selected from literature for development of look-up table.	J.29